arXiv:1512.02364v1 [math.CA] 8 Dec 2015

Concavity of some entropies

Ioan RAŞA
Department of Mathematics, Technical University of Cluj-Napoca, Romania
Email address: Ioan.Rasa@math.utcluj.ro

Abstract

It is well-known that the Shannon entropies of some parameterized probability distributions are concave functions with respect to the parameter. In this paper we consider a family of such distributions (including the binomial, Poisson, and negative binomial distributions) and investigate the concavity of the Shannon, Rényi, and Tsallis entropies of them.

1 Introduction

Let $c \in \mathbb{R}$, $I_c := \left[0, -\frac{1}{c}\right]$ if $c < 0$, and $I_c := [0, +\infty)$ if $c \geq 0$.

For $\alpha \in \mathbb{R}$ and $k \in \mathbb{N}_0$ the binomial coefficients are defined as usual by

$$\binom{\alpha}{k} := \frac{\alpha(\alpha-1)\dots(\alpha-k+1)}{k!} \quad \text{if } k \in \mathbb{N}, \text{ and } \binom{\alpha}{0} := 1.$$

Let $n > 0$ be a real number such that $n > c$ if $c \geq 0$, or $n = -cl$ with some $l \in \mathbb{N}$ if $c < 0$.

For $k \in \mathbb{N}_0$ and $x \in I_c$ define

$$p_{n,k}^{[c]}(x) := (-1)^k \binom{-\frac{n}{c}}{k} (cx)^k (1+cx)^{-\frac{n}{c}-k}, \quad \text{if } c \neq 0,$$

$$p_{n,k}^{[0]}(x) := \lim_{c \to 0} p_{n,k}^{[c]}(x) = \frac{(nx)^k}{k!} e^{-nx}.$$

Details and historical notes concerning these functions can be found in [3], [7], [21] and the references therein. In particular,

$$\frac{d}{dx} p_{n,k}^{[c]}(x) = n \left(p_{n+c,k-1}^{[c]}(x) - p_{n+c,k}^{[c]}(x) \right). \tag{1.1}$$

Moreover, $\sum_{k=0}^{\infty} p_{n,k}^{[c]}(x) = 1$, so that $\left(p_{n,k}^{[c]}(x)\right)_{k \geq 0}$ is a parameterized probability distribution. Its associated Shannon entropy is

$$H_{n,c}(x) := -\sum_{k=0}^{\infty} p_{n,k}^{[c]}(x) \log p_{n,k}^{[c]}(x),$$

while the Rényi entropy of order 2 and the Tsallis entropy of order 2 are given, respectively, by (see [18], [20])

$$R_{n,c}(x) := -\log S_{n,c}(x); \quad T_{n,c}(x) := 1 - S_{n,c}(x),$$

where

$$S_{n,c}(x) := \sum_{k=0}^{\infty} \left(p_{n,k}^{[c]}(x)\right)^2, \quad x \in I_c.$$

The cases $c = -1$, $c = 0$, $c = 1$ correspond, respectively, to the binomial, Poisson, and negative binomial distributions.

2 Shannon entropy

$H_{n,-1}$ is a concave function; this is a special case of the results of [19]; see also [6], [8], [9] and the references therein. $H_{n,0}$ is also concave; moreover, $H'_{n,0}$ is completely monotonic (see, e.g., [2, p. 2305]).

For the sake of completeness we present here the proof for the concavity of $H_{n,c}$, $c \in \mathbb{R}$. Let us consider separately the cases $c \geq 0$ and $c < 0$.

Theorem 2.1. *For $c \geq 0$, $H_{n,c}$ is concave and increasing on $[0, +\infty)$.*

Proof. Using (1.1), it is a matter of calculus to prove that

$$H'_{n,c}(x) = n \sum_{k=0}^{\infty} p_{n+c,k}^{[c]}(x) \log \frac{p_{n,k}^{[c]}(x)}{p_{n,k+1}^{[c]}(x)},$$

which leads to

$$H'_{n,c}(x) = n \left(\log \frac{1+cx}{x} + \sum_{k=0}^{\infty} p_{n+c,k}^{[c]}(x) \log \frac{k+1}{n+ck} \right), \tag{2.1}$$

and therefore to

$$H''_{n,c}(x) = -\frac{n}{x(1+cx)} + n(n+c) \sum_{k=0}^{\infty} p_{n+2c,k}^{[c]}(x) \log \frac{(k+2)(n+ck)}{(k+1)(n+ck+c)}.$$

It follows that

$$H''_{n,c}(x) > -\frac{n}{x(1+cx)}, \quad x > 0. \tag{2.2}$$

Since $\log t < t - 1$, $t > 1$, we have also

$$\begin{aligned} H''_{n,c}(x) &< -\frac{n}{x(1+cx)} + n(n+c) \sum_{k=0}^{\infty} p_{n+2c,k}^{[c]}(x) \frac{n-c}{(k+1)(n+ck+c)} \\ &= -\frac{n}{x(1+cx)} \left(\frac{c}{n} + \left(1 - \frac{c}{n}\right) (1+cx)^{-\frac{n}{c}} \right) < 0, \end{aligned}$$

so that $H_{n,c}$ is concave on $[0, +\infty)$; being positive, it is also increasing on $[0, +\infty)$. □

Remark 2.2. *From* (2.2) *it follows that the functions* $H_{n,0}(x)+nx\log x$ *and* $H_{n,c}(x)+\frac{n}{c}\left(cx\log x-(1+cx)\log(1+cx)\right)$, $c>0$, *are convex on* $[0,+\infty)$.

Remark 2.3. *The following inequalities are valid for* $x>0$ *and* $c\geq 0$*:*

$$\log\frac{x}{1+cx}\leq\sum_{k=0}^{\infty}p_{n+c,k}^{[c]}(x)\log\frac{k+1}{ck+n}\leq\log\frac{x+\frac{1}{n}}{1+cx}. \tag{2.3}$$

The first one follows from $H'_{n,0}(x)>0$, taking into account (2.1), and the second is a consequence of Jensen's inequality applied to the concave function $\log x$. In particular, for $c=0$ and $n=1$ we get:

$$\log x\leq\sum_{k=0}^{\infty}e^{-x}\frac{x^k}{k!}\log(k+1)\leq\log(x+1),\quad x>0.$$

The case $c<0$ can be studied with the same method as in Theorem 2.1, but we present here a different approach, based on an integral representation from [10].

Theorem 2.4. *For* $c<0$, $H_{n,c}$ *is concave on* $\left[0,-\frac{1}{c}\right]$, *increasing on* $\left[0,-\frac{1}{2c}\right]$, *and decreasing on* $\left[-\frac{1}{2c},-\frac{1}{c}\right]$.

Proof. For $c<0$ we have $n=-cl$ with $l\in\mathbb{N}$. Using [10, (2.5)] we get

$$H_{n,c}(x)=H_{l,-1}(-cx)=-l\left[(-cx)\log(-cx)+(1+cx)\log(1+cx)\right]+$$
$$\int_0^1\frac{s-1}{\log s}\frac{(1+cx-cxs)^l+\left((1+cx)s-cx\right)^l-1-s^l}{(s-1)^2}ds.$$

It is matter of calculus to prove that

$$H''_{n,c}(x)=-\frac{n}{x(1+cx)}+c^2l(l-1)\int_0^1\frac{s-1}{\log s}\left[(1+cx-cxs)^{l-2}+\left((1+cx)\,s-cx\right)^{l-2}\right]ds. \tag{2.4}$$

For $0<s<1$ we have $0<\frac{s-1}{\log s}<1$; moreover

$$\int_0^1\left[(1+cx-cxs)^{l-2}+\left((1+cx)\,s-cx\right)^{l-2}\right]ds=\frac{1}{l-1}\left[\frac{1-(-cx)^{l-1}}{1+cx}-\frac{1-(1+cx)^{l-1}}{cx}\right].$$

Summing up, we get

$$H''_{n,c}(x)<-n\frac{(1+cx)^{-\frac{n}{c}}+(-cx)^{-\frac{n}{c}}}{x(1+cx)}<0,\quad 0<x<-\frac{1}{c}.$$

Consequently, $H_{n,c}$ is concave on $\left[0,-\frac{1}{c}\right]$. Since

$$H_{n,c}\left(-\frac{1}{2c}-t\right)=H_{n,c}\left(-\frac{1}{2c}+t\right),\quad t\in\left[0,-\frac{1}{2c}\right],$$

we conclude that $H_{n,c}$ is increasing on $\left[0,-\frac{1}{2c}\right]$ and decreasing on $\left[-\frac{1}{2c},-\frac{1}{c}\right]$. □

Remark 2.5. *Let $c < 0$ and $l \geq 2$. From (2.4) it follows that $H''_{n,c}(x) > -\frac{n}{x(1+cx)}$, and so the function*

$$H_{n,c}(x) + \frac{n}{c}\left[cx\log x - (1+cx)\log(1+cx)\right]$$

is convex on $\left[0, -\frac{1}{c}\right]$.

Remark 2.6. *For $c = -1$, the method used to prove (2.3) leads to*

$$\begin{aligned}\log\frac{x}{1-x} &< \sum_{k=0}^{n}\binom{n}{k}x^k(1-x)^{n-k}\log\frac{k+1}{n+1-k}\\ &< \log\left(\frac{x}{1-x}+\frac{1-(n+2)x^{n+1}}{(n+1)(1-x)}\right), \quad 0<x<\frac{1}{2}.\end{aligned}$$

3 $S_{n,c}$ and Heun functions

The following conjecture was formulated in [13]:

Conjecture 3.1. *$S_{n,-1}$ is convex on $[0,1]$.*

Th. Neuschel [11] proved that $S_{n,-1}$ is decreasing on $\left[0, \frac{1}{2}\right]$ and increasing on $\left[\frac{1}{2}, 1\right]$. The conjecture and the result of Neuschel can be found also in [5].

A proof of the conjecture was given by G. Nikolov [12], who related it with some new inequalities involving Legendre polynomials. Another proof can be found in [4].

Using the important results of Elena Berdysheva [3], the following extension was obtained in [17]:

Theorem 3.2. *([17, Theorem 9]). For $c < 0$, $S_{n,c}$ is convex on $\left[0, -\frac{1}{c}\right]$.*

A stronger conjecture was formulated in [14] and [17]:

Conjecture 3.3. *For $c \in \mathbb{R}$, $S_{n,c}$ is logarithmically convex, i.e., $\log S_{n,c}$ is convex.*

It was validated for $c \geq 0$ by U. Abel, W. Gawronski and Th. Neuschel [1], who proved a stronger result:

Theorem 3.4. *([1]). For $c \geq 0$, the function $S_{n,c}$ is completely monotonic, i.e.,*

$$(-1)^m\left(\frac{d}{dx}\right)^m S_{n,c}(x) > 0, \quad x \geq 0, m \geq 0.$$

Consequently, for $c \geq 0$, $S_{n,c}$ is logarithmically convex, and hence convex.

On the other hand, according to [17, Th. 4], $S_{n,c}$ is a solution to the differential equation

$$\begin{aligned}x(1+cx)(1+2cx)y''(x) + (4(n+c)x(1+cx)+1)\,y'(x) +\\ +2n(1+2cx)y(x) = 0.\end{aligned} \tag{3.1}$$

Consequently, for $c \neq 0$ the function $S_{n,c}\left(-\frac{x}{c}\right)$ is a solution to the Heun equation

$$y''(x) + \left(\frac{1}{x} + \frac{1}{x-1} + \frac{\frac{2n}{c}}{x-\frac{1}{2}}\right) y'(x) + \frac{\frac{2n}{c}x - \frac{n}{c}}{x(x-1)\left(x-\frac{1}{2}\right)} y(x) = 0,$$

and $S_{n,0}$ is a solution of the confluent Heun equation:

$$u''(x) + \left(4n + \frac{1}{x}\right) u'(x) + \frac{2nx - 2n}{x(x-1)} u(x) = 0.$$

For details, see [14]-[17].

4 Rényi entropy and Tsallis entropy

Theorem 4.1. *(i) For $c \geq 0$, $R_{n,c}$ is concave and increasing on $[0, +\infty)$.*

(ii) $R'_{n,c}$, with $c \in \mathbb{R}$, is a solution to the Riccati equation

$$\begin{aligned} x(1+cx)(1+2cx)u'(x) = x(1+cx)(1+2cx)u^2(x) - \\ - \left(4(n+c)x(1+cx)+1\right) u(x) + 2n(1+2cx). \end{aligned}$$

Proof. i) is a direct consequence of Theorem 3.4.

ii) We have $S_{n,c} = \exp(-R_{n,c})$ and (3.1) yields

$$\begin{aligned} x(1+cx)(1+2cx)\left((R'_{n,c})^2 - R''_{n,c}\right) - \\ - \left(4(n+c)x(1+cx)+1\right) R'_{n,c} + 2n(1+2cx) = 0. \end{aligned}$$

Setting $u = R'_{n,c}$, we conclude the proof.

□

Remark 4.2. *As far as we know, Conjecture 3.3 is still open for $c < 0$, so that the concavity of $R_{n,c}$, $c < 0$, remains to be investigated.*

Theorem 4.3. *(i) $T_{n,c}$ is concave. For $c \geq 0$ it is increasing on $[0, +\infty)$. For $c < 0$ it is increasing on $\left[0, -\frac{1}{2c}\right]$ and decreasing on $\left[-\frac{1}{2c}, -\frac{1}{c}\right]$.*

(ii) $T_{n,c}$ is a solution to the equation

$$\begin{aligned} x(1+cx)(1+2cx)u''(x) + \left(4(n+c)x(1+cx)+1\right) u'(x) + \\ + 2n(1+2cx)u(x) = 2n(1+2cx). \end{aligned}$$

Proof. (i) is a consequence of Theorems 3.2 and 3.4, while

(ii) follows from (3.1).

□

5 Some inequalities

a) The explicit expression of $S_{n,-1}$, $n \in \mathbb{N}$, is

$$S_{n,-1}(x) = \sum_{k=0}^{n} \left(\binom{n}{k} x^k (1-x)^{n-k} \right)^2, \quad x \in [0,1].$$

Consider also the function

$$f_n(t) := \sum_{j=0}^{n-1} \binom{n-1}{j} \left(\binom{n}{j+1} t^{2j+1} - \binom{n}{j} t^{2j} \right), \quad t \geq 1.$$

Since $S_{n,-1}(1-x) = S_{n,-1}(x)$, it follows that

$$S_{n,-1}^{(2j+1)}\left(\frac{1}{2}\right) = 0, \quad j = 0, 1, \ldots, n-1.$$

In relation with Conjecture 3.1, it was also conjectured in [13] that

$$S_{n,-1}^{(2j)}\left(\frac{1}{2}\right) > 0, \quad j = 0, 1, \ldots, n, \tag{5.1}$$

$$f_n^{(i)}(t) \geq 0, \quad t \geq 1; \quad i = 0, 1, \ldots, 2n-1, \tag{5.2}$$

$$\sum_{j=[i/2]}^{n-1} \binom{n-1}{j} \left(\binom{n}{j+1} \binom{2j+1}{i} - \binom{n}{j} \binom{2j}{i} \right) \geq 0, \quad i = 0, 1, \ldots, 2n-1. \tag{5.3}$$

We shall prove here these inequalities[1].

It can be proved directly that

$$S_{n,-1}\left(\frac{1}{2}\right) = \frac{1}{4^n}\binom{2n}{n}; \quad S_{n,-1}^{(2)}\left(\frac{1}{2}\right) = \frac{1}{4^{n-2}}\binom{2n-2}{n-1};$$

$$S_{n,-1}^{(4)}\left(\frac{1}{2}\right) = \frac{9}{4^{n-4}}\binom{2n-4}{n-2}.$$

The following formula was obtained in [4]:

$$S_{n,-1}(x) = \sum_{k=0}^{n} 4^{k-n} \binom{2n}{n} \binom{n}{k}^2 \binom{2n}{2k}^{-1} \left(x - \frac{1}{2}\right)^{2k}. \tag{5.4}$$

[1] For (5.2) with $i = 0$, see also
http://www.artofproblemsolving.com/community/c6h494060p2772738.

Using it we get

$$S_{n,-1}^{(2j)}\left(\frac{1}{2}\right) = (2j)!4^{j-n}\binom{2n}{n}\binom{n}{j}^2\binom{2n}{2j}^{-1}, \quad j = 0, 1, \ldots, n,$$

and so (5.1) is proved.

On the other hand, it is not difficult to prove that

$$f_n(t) = \frac{1}{2n}(t+1)^{2n-1}S'_{n,-1}\left(\frac{t}{t+1}\right), \quad t \geq 1. \tag{5.5}$$

From (5.4) we obtain

$$S'_{n,-1}(x) = \sum_{k=1}^{n} 2ka_{nk}\left(x - \frac{1}{2}\right)^{2k-1} \tag{5.6}$$

for certain $a_{nk} > 0$. Now (5.5) and (5.6) imply

$$f_n(t) = \sum_{k=1}^{n} c_{nk}(t-1)^{2k-1}(t+1)^{2n-2k}, \quad t \geq 1,$$

with suitable $c_{nk} > 0$, and using Leibniz' formula we get (5.2).

Finally, starting from the definition of $f_n(t)$, it is not difficult to infer that

$$f_n^{(i)}(1) = i! \sum_{j=[i/2]}^{n-1} \binom{n-1}{j}\left(\binom{n}{j+1}\binom{2j+1}{i} - \binom{n}{j}\binom{2j}{i}\right).$$

Combined with (5.2), this proves (5.3).

b) Let $B_m : \mathscr{C}[0,1] \to \mathscr{C}[0,1]$, $B_m f(x) = \sum_{k=0}^{m}\binom{m}{k}x^k(1-x)^{m-k}$, be the classical Bernstein operators. It is well-known that if $f \in \mathscr{C}[0,1]$ is convex, then $B_m f$ is also convex.

Now let $w_n \in \mathscr{C}[0,1]$ be piece-wise linear, such that $w_n\left(\frac{2k-1}{2n}\right) = 0$, $k = 1, \ldots, n$; $w_n\left(\frac{2k}{2n}\right) = \binom{n}{k}^2/\binom{2n}{2k}$, $k = 0, 1, \ldots, n$. Then $B_{2n}w_n = S_{n,-1}$, hence $B_{2n}w_n$ is convex although the graph of w_n is "like a saw".

References

[1] U. Abel, W. Gawronski, Th. Neuschel, Complete monotonicity and zeros of sums of squared Baskakov functions, Appl. Math. Comput. 258 (2015), 130-137.

[2] J.A. Adell, A. Lekuona and Y. Yu, Sharp bounds on the entropy of the Poisson Law and related quantities, IEEE Trans. Information Theory 56 (2010), 2299-2306.

[3] E. Berdysheva, Studying Baskakov-Durrmeyer operators and quasi-interpolants via special functions, J. Approx. Theory 149 (2007), 131-150.

[4] I. Gavrea, M. Ivan, On a conjecture concerning the sum of the squared Bernstein polynomials, Appl. Math. Comput. 241 (2014), 70-74.

[5] H. Gonska, I. Raşa, M.-D. Rusu, Chebyshev-Grüss-type inequalities via discrete oscillations, Bul. Acad. Stiinte Repub. Mold. Mat. no. 1 (74) (2014), 63-89; arxiv 1401.7908 [math.CA].

[6] P. Harremoës, Binomial and Poisson distributions as maximum entropy distributions, IEEE Trans. Information Theory 47 (2001), 2039 - 2041.

[7] M. Heilmann, Erhöhung der Konvergenzgeschwindigkeit bei der Approximation von Funktionen mit Hilfe von Linearkombinationen spezieller positiver linearer Operatoren, Habilitationsschrift, Universität Dortmund (1992).

[8] E. Hillion, Concavity of entropy along binomial convolutions, Electron. Commun. Probab. 17 (2012), 1-9.

[9] E. Hillon, O. Johnson, A proof of the Shepp-Olkin entropy concavity conjecture, arXiv: 1503.01570v1, 5 Mar 2015.

[10] C. Knessl, Integral representations and asymptotic expansions for Shannon and Rényi entropies, Appl. Math. Lett. 11 (1998), 69-74.

[11] Th. Neuschel, Unpublished manuscript (2012).

[12] G. Nikolov, Inequalities for ultraspherical polynomials. Proof of a conjecture of I. Raşa, J. Math. Anal. Appl. 418 (2014), 852-860.

[13] I. Raşa, Unpublished manuscripts (2012).

[14] I. Raşa, Special functions associated with positive linear operators, arxiv: 1409.1015v2.

[15] I. Raşa, Rényi entropy and Tsallis entropy associated with positive linear operators, arxiv: 1412.4971v1.

[16] I. Raşa, Entropies and the derivatives of some Heun functions, arxiv: 1502.05570v1.

[17] I. Raşa, Entropies and Heun functions associated with positive linear operators, Appl. Math. Comput. 268 (2015), 422-431.

[18] A. Rényi, On measures of entropy and information, Proc. Fourth Berkeley Symp. Math. Statist. Prob., Vol. 1, Univ. of California Press (1961), pp. 547-561.

[19] L.A. Shepp, I. Olkin, Entropy of the sum of independent Bernoulli random variables and of the multinomial distribution, in: Proc. Contributions to Probability, New York (1981), pp. 201-206.

[20] C. Tsallis, Possible generalization of Boltzmann-Gibbs statistics, J. Stat. Phys. 52 (1988), 479-487.

[21] M. Wagner, Quasi-Interpolaten zu genuinen Baskakov-Durrmeyer-Typ Operatoren, Shaker Verlag, Aachen (2013).